



FIFRA 2(ee) RECOMMENDATION
Expires December 31, 2005

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF TEXAS

Imidan[®] 70-W
Agricultural Insecticide
EPA Reg. No. 10163-169

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

This recommendation is made as permitted under FIFRA section 2(ee) and has not been submitted to or approved by the EPA.

CROP RECOMMENDATIONS

CROP	PEST CONTROLLED	USE RATE OF IMIDAN 70-WP OR IMIDAN 70-WSB PER ACRE
Pecans	Nut Casebearer	1 to 1½ lbs.
	Hickory Shuckworm	1½ to 2 lbs.
	Pecan Weevil*	2 to 3 ¹ / ₈ lbs.
	Black Pecan Aphid	2 lbs.

*For control of low to moderate populations of Pecan Weevil; adequate control may not be achieved when heavy populations are present.

- Use the highest rate listed when insect pressure is severe. Thorough coverage is necessary for effective control.
- Repeat applications as necessary based on insect infestations or local or State programs.
- The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum insecticidal activity. For maximum performance, sprays with Imidan should have a pH of 5 to 6.

RESTRICTIONS

Do not exceed 3¹/₈ lbs. per acre per application.

Do not apply within 14 days of harvest.

Do not graze livestock on cover crops grown in treated pecan groves.