



FIFRA §2(ee) RECOMMENDATION  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA,  
ARKANSAS, FLORIDA, GEORGIA, KANSAS, KENTUCKY, LOUISIANA,  
MISSISSIPPI, OKLAHOMA, TENNESSEE

# **Imidan<sup>®</sup> 70-W**

*Agricultural Insecticide*

EPA Reg. No. 10163-169

## DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

## FOR CONTROL OF NUT CASEBEARER, HICKORY SHUCKWORM, PECAN WEEVIL\*, AND BLACK PECAN APHID ON PECANS

CROP	PEST	USE RATE OF IMIDAN 70-W PER ACRE
PECANS	Nut Casebearer	1 to 1½ lbs.
	Hickory Shuckworm	1½ to 2 lbs.
	Pecan Weevil*	2 to 3 <sup>1</sup> / <sub>8</sub> lbs.
	Black Pecan Aphid	2 lbs.

\*For control of low to moderate populations of Pecan Weevil; adequate control may not be achieved when heavy populations are present.

## RECOMMENDATIONS:

- Use the highest rate listed when insect pressure is severe. Thorough coverage is necessary for effective control.
- Repeat applications as necessary based on insect infestations or local or State programs.
- The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum insecticidal activity. For maximum performance, sprays with Imidan should have a pH of 5 to 6.

## RESTRICTIONS:

- Do not exceed 3<sup>1</sup>/<sub>8</sub> lbs. per acre per application.
- Do not apply within 14 days of harvest.
- Do not graze or feed livestock on cover crops grown in treated pecan groves.

For additional information, contact your regional Gowan Company sales representative or your local agrichemical dealer.

Imidan<sup>®</sup> is a registered trademark of Gowan Company, L.L.C.

99-02-02