

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA**



EPA REG. NO: 10163-357

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered labels are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

PRODUCT INFORMATION

EcoSwing is a biofungicide for use on growing crops in agricultural, greenhouse, ornamental and turf use sites to control fungal diseases such as powdery mildew (oidiums), sour rot, and *Botrytis cinerea*. Optimum disease control is achieved when the biofungicide is used according to label directions and applied in a regularly scheduled preventative spray program.

The use of EcoSwing is compatible with the principles of Integrated Pest Management (IPM) and programs that attempt to minimize disease resistance to fungicides. Unlike single-site mode of action fungicides which are at risk from disease resistance, EcoSwing has a multi-site mode of action, and may be used to delay or prevent the development of resistance to single site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of EcoSwing in IPM and resistance management programs.

PRE-HARVEST INTERVAL: EcoSwing has a zero day pre-harvest interval and can be applied up to and including the day of harvest

APPLICATION DIRECTIONS

CROP	PEST	FIELD RATE	COMMENTS
Snap Beans	White mold (<i>Sclerotinia sclerotiorum</i>)	2 pts/Acre	For best results, apply at least every 7-14 days on a regular basis before the level of infestation becomes heavy.

EcoSwing™ is a trademark used under license by Gowan Company, L.L.C.