

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF IDAHO For Control of Powdery Mildew (*Erysiphe heraclei*) in Carrots Grown for Seed

TETRACONAZOLE GROUP 3 FUNGICIDE



EPA Reg. No 10163-397 / EPA SLN NO. ID-220006

This label is valid until December 31, 2026, or until otherwise amended, withdrawn, canceled, or suspended.

 Active Ingredient:
 20.5%

 Tetraconazole*
 20.5%

 Other Ingredients:
 79.5%

 Total
 100.0%

DOMARK 230 ME FUNGICIDE is a micro emulsion containing 1.9 pounds of Tetraconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2,-tetraflurorethoxy)propyl]1H-1,2,4-triazole

- The Section 3 product label and this SLN Supplemental label must be in the possession of the user at the time
 of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered Domark 230 ME label.
- Use of Domark 230 ME according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Domark 230ME.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise caution when making applications of DOMARK 230 ME FUNGICIDE, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment wash waters or rinsate.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all activities with the exception of 3 days for detasseling corn grown for seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any water proof material
- Shoes plus socks

CROP	RATE	TARGET DISEASE	COMMENTS	
Carrots Grown for Seed	6.7 fl ounces / acre (0.1 lb. ai / A)	Powdery Mildew (<i>Erysiphe heraclei</i>)	 20 gallons of water minimum per acre for ground application or in a minimum of 10 gallons of water per acre for aerial application. Begin applications prior to or at the start of disease development and repeat if needed 14 to 21 days after the first application. 	
	RESTRICTIONS / PRECAUTIONS:			
	Do not apply more than 13.4 fl oz of product per acre per year.			
	 Do not ap 	Do not apply more than 2 applications of Domark 230 ME fungicide per acre per year		
	Do not apply within 14 days between multiple applications.			
	 No portion of the carrot seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, leaves and seed screenings may be used or distributed for food or feed purposes. Carrot seed may not be distributed for human consumption or animal feed. 			

Distributed by: Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569