



**2(ee) Recommendation
EPA Reg. No. 10163-332**

For use in the States of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, New York, North Dakota, Ohio, Pennsylvania, South Dakota, and Wisconsin.

Fungicide for Control of Tar Spot on Field Corn, Popcorn, Corn Grown For Seed Production.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) and the entire label on the product container before using this product.

Field Corn, Popcorn, Corn Grown For Seed Production

Target Disease	Product Use Rate per Application fl oz/A	Use Directions
Tar Spot <i>(Phyllachora maydis)</i>	10.0-17.0	Apply prior to disease onset when conditions favor disease development. A second application may be made no fewer than 7 days later but not to exceed 17.06 fl oz per acre per year. Use AFFIANCE® Fungicide as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants.

RESTRICTIONS AND LIMITATIONS

1. Do not make more than two (2) applications per year.
2. Do not make more than 2 sequential applications before alternating to another fungicide with a different mode of action.
3. Do not apply more than 17.06 fl oz of AFFIANCE® Fungicide per acre per year.
4. Do not apply more than 0.09 lb ai of a tetraconazole-containing product per acre per year.
5. Do not apply more than 2.0 lb ai of an azoxystrobin-containing product per acre per year.
6. Do not apply AFFIANCE® Fungicide after corn growth stage R3 (brown silk/milk).
7. Do not use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. A compatibility agent, another fungicide, or an insecticide may be included if needed and labeled for use in corn. Refer to adjuvant product label for specific use directions and restrictions. Always follow the more restrictive label.
8. For chemigation, apply in 0.1 – 0.25 inches/A of water.
9. Do not apply within 7 days of harvest (7 day PHI).
10. Do not harvest silage within 21 days of an application.

IMPORTANT NOTICE

Follow all applicable directions, restrictions and precautions.

This bulletin contains supplemental instructions for use of the referenced **AFFIANCE®** product. Follow these instructions carefully. This bulletin must be in the possession of the user at the time of application. Read the entire label before using this product, including the "LIMITATION OF WARRANTY AND LIABILITY" in the label for **AFFIANCE®** before using.

ID# 19-03-02

AFFIANCE® is a registered trademark of Gowan Company, LLC

