

**FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY
WITHIN THE STATE OF TEXAS**



**EPA REG. NO. 10163-397
Expiration Date: 12-31-2026**

Fungicide for Control of Listed Diseases Peanuts

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Peanut

Target Disease	Product Use Rate per Application fl oz/A	Use Directions
<p>Early leaf Spot (Cercospora arachidicola)</p> <p>Late leaf spot (Cercosporidium personatum)</p>	<p>2.5 – 5.25</p> <p>(0.037 – 0.078 lb. ai/A)</p>	<p>Begin applications prior to onset of disease when conditions are favorable for disease development, generally around 30 to 40 days after planting.</p> <p>Reapply DOMARK 230 ME using a 14 day interval.</p> <p>DOMARK 230 ME may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.</p> <p>Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results.</p> <p>Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.</p>



Produced For:
Gowan Company LLC
P.O. BOX 5569
Yuma, AZ 85366-5569