

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA AND CALIFORNIA**



EPA Reg. No. 10163-200

For use on Bulb Onions for suppression of Crabgrass (*Digitaria spp.*), Fall Panicum (*Panicum dichotomiflorum*), Foxtail (*Setaria spp.*), Goosegrass (*Eleusine indica*), Jungle Rice (*Echinochloa colonum*), Sprangletop (*Leptochloa spp.*), Barnyardgrass (*Echinochloa crusgalli*), Lambsquarters (*Chenopodium album*), Purslane (*Portulaca oleracea*), Redroot Pigweed (*Amaranthus retroflexus*), Burning Nettle (*Urtica urens*), Nettleleaf Goosefoot (*Chenopodium murale*)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

Crop	Weeds Suppressed	BROADCAST RATE (Quarts/Acre)	COMMENTS
Bulb Onions	Crabgrass (<i>Digitaria spp.</i>) Fall Panicum (<i>Panicum dichotomiflorum</i>) Foxtail (<i>Setaria spp.</i>) Goosegrass (<i>Eleusine indica</i>) Jungle Rice (<i>Echinochloa colonum</i>) Sprangletop (<i>Leptochloa spp.</i>) Barnyardgrass (<i>Echinochloa crusgalli</i>) Lambsquarters (<i>Chenopodium album</i>) Purslane (<i>Portulaca oleracea</i>) Redroot Pigweed (<i>Amaranthus retroflexus</i>) Burning Nettle (<i>Urtica urens</i>) Nettleleaf Goosefoot (<i>Chenopodium murale</i>)	2 - 3 (2 - 3 lb a.i./acre)	Do not apply more than 6 lbs. a.i. per acre per season. Apply preplant or preemergence. Follow application instructions given on this label. Onions germinate and grow best when planted in fine-textured soil with good water drainage where the soil salinity value is less than 4 ECe* and soil temperature is between 55° and 75°. Use of Prefar on onions under less favorable soil and environmental conditions may reduce stand and vigor resulting in lower yield. To avoid injury, do not incorporate by mechanical methods in California or Arizona. Consult your local extension agent or university specialist for further information. (*ECe is the electrical conductivity extract.) For suppression of redroot pigweed and common lambsquarters, prepare a good, flat seed bed by standard cultural practices. Apply Prefar 4-E in the fall in 10 - 50 gals. of water in a 10 - 12 inch band and bed-up, throwing soil from the furrows over the sprayed band of Prefar 4-E. Allow to remain undisturbed until spring. Prior to planting in spring, drag-off the bed tops being careful not to drag off soil below the level of Prefar 4-E application. Plant in the center of the beds.

Combining Prefar 4-E with fertilizers or other agricultural chemicals could result in incompatibility, reduced efficacy, or reduced crop safety. It is recommended that the user test the response of combination treatments with other products on a small scale prior to commercial scale use.

Prefar® 4-E is a registered trademark of Gowan Company, L.L.C.



Produced for:
Gowan Company, LLC
P.O. Box 5569
Yuma, AZ 85366-5569