

**FIFRA 2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE IN FLORIDA ONLY
SUPPLEMENTAL LABELING FOR CONTROL OF ADDITIONAL PESTS IN
PEPPERS**

Lorsban* 75WG

EPA Reg. No. 62719-301-10163

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA

Crop	Pest	Rate (lbs./acre)
Peppers	Pepper Weevil Western Flower Thrips	1.33 lbs

DIRECTIONS FOR USE

Use and Dosage Recommendations

Apply 1 1/3 lb of Lorsban 75WG (1 1.33-lb packet) in 50 to 150 gallons of water per acre as a broadcast foliar spray using power-operated ground application equipment. Apply when field counts indicate damaging populations are developing or present. To avoid possible phytotoxicity, tank mixes must be tested on a small area prior to use on the remaining acres.

Specific Use Restrictions

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not make more than 8 applications of products containing chlorpyrifos per crop.
- Maximum single application rate is 1.33 lb of Lorsban 75WG (1 lb ai chlorpyrifos) per acre.
- **Minimum Re-treatment interval:** Do not make a second application of Lorsban 75WG or other product containing chlorpyrifos within 10 days of the first application.
- **Chemigation:** Do not apply Lorsban 75WG through any type of irrigation system.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not apply Lorsban 75WG within 10 days of transplanting or to plants that are under extreme heat or drought stress.

*Trademark of Dow AgroSciences LLC