

**This supplemental label expires on March 24, 2019
and must not be used or distributed after this date.**

Onager[®]

Miticide

EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.8%
OTHER INGREDIENTS	<u>88.2%</u>
TOTAL	100.00%

Onager contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

CROP	PEST	RATE Oz./acre	COMMENTS
COTTON (35 day PHI)	Twospotted spider mite, Pacific spider mite, Strawberry spider mite, Carmine spider mite	12 - 20	Apply Onager before boll opening and prior to adult mite buildup. Complete coverage of foliage is essential to optimal performance. Ground Application: Apply at recommended rates Air application: Use recommended per acre rate and apply a minimum of 2 gallons of finished spray per acre. Chemigation: see details under Chemigation Systems.
	<ul style="list-style-type: none"> • Do not apply more than a total of 20 oz. formulated product per acre per year. • Do not make more than one application of Onager or any other hexythiazox product to the same crop per year. 		

NET CONTENTS _____



EPA Reg. No. 10163-277
EPA Est. No.

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Refer to the Section 3 label for the use.
EPA Text: Onager (EPA 3-24-16)