

**FIFRA 2(ee) RECOMMENDATION  
FOR DISTRIBUTION AND USE IN MICHIGAN ONLY  
SUPPLEMENTAL LABELING FOR CONTROL OF ADDITIONAL PEST IN SOUR  
CHERRIES**

# Lorsban<sup>®</sup> 75WG

**EPA Reg. No. 62719-301-10163**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA

**TREE FRUITS AND TREE NUTS (FOLIAR SPRAYS)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days for tree fruits and 24 hours for tree nuts unless PPE required for early entry is worn.

Apply Lorsban 75WG as a foliar spray at the dosages indicated to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop and apply as a concentrate or dilute spray using conventional, power-operated spray equipment. For dilute sprays applied to tree nut crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of Lorsban 75WG per acre. Treat when pests appear or in accordance with local conditions. Aerial application may result in less effective insect control because of reduced coverage. Consult your State agricultural experiment station, certified pest control advisor, or extension service specialist for specific use information in your area.

Crop	Pest	Rate (lbs./acre)
Cherry (sour)	Plum Curculio	1.33 - 2

**Specific Use Precautions:**

- Lorsban 75WG is highly toxic to bees exposed to direct treatment and must not be applied when bees are actively foraging in the treated area.
- To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours of application of Lorsban 75WG.

**Specific Use Restrictions:**

Crop	Preharvest Interval (Days)	Total Number of Sprays	Total Amount Chlorpyrifos per Season (lb ai/acre)	Total Amount Lorsban 75WG per Season (lb/acre)
Sour Cherry	14	8	12	16

- Do not make a second application of Lorsban 75WG or other product containing chlorpyrifos within 10 days of the first application.
- Do not allow meat or dairy animals to graze in treated orchards.