

Onager[®]

Miticide

EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.8%
OTHER INGREDIENTS	88.2%
TOTAL	100.00%

Onager contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

CROP	PEST	RATE Oz./acre	COMMENTS
Wheat, including emmer wheat and triticale (West of the Rockies)	Banks grass mite	10 - 24	Apply at the first sign of mites before populations begin to build. Ground Application: Apply the recommended rate of Onager by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems on Section 3 label.
			<ul style="list-style-type: none"> • Do not apply after the boot stage of growth. • Do not harvest hay or forage for at least 14 days after application. • Do not apply more than a total of 24 oz. formulated product per acre per year. • Do not make more than one application per year.

NET CONTENTS _____



EPA Reg. No. 10163-277
EPA Est. No.

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

This supplemental expires on 12-31-18. Refer to the Section 3 label for the use.