

**REGISTRATION FOR SPECIAL LOCAL NEED  
FOR DISTRIBUTION AND USE ONLY WITHIN COLORADO**



**EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY**

EPA Reg. No. 10163-277

This label expires and must not be distributed or used after December 31, 2021.

<b>ACTIVE INGREDIENT:</b> Hexythiazox	<b>% By Wt.</b>
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide .....	11.8%
<b>OTHER INGREDIENTS</b> .....	<b>88.2%</b>
	<b>TOTAL 100.00%</b>

Onager contains 1.0 lb. active ingredient per gallon  
Contains Petroleum Distillate

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, Worker Protection Standard Requirements and precautions on the USEPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

CROP	PEST	RATE OzS./Acre	COMMENTS
GRASS SEED	Banks grass mite, Twospotted Spider Mite	12 – 24	<p><b>Ground Application:</b> Apply the recommended rate of Onager by ground equipment at 15 – 20 GPA.</p> <p><b>Air Application:</b> Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p><b>Chemigation:</b> This product may be applied through irrigation systems*. Refer to the EPA registered label for chemigation directions.</p>
	<ul style="list-style-type: none"> <li>• Do not cut current year's treated seed crop for hay or forage.</li> <li>• Do not graze or feed livestock on cover crops growing in treated areas.</li> <li>• Screenings from seed processing are prohibited from food or feed channels. All Onager treated seed screenings must be removed from the food or feed market.</li> <li>• Treated seed is not to be used for sprouting. All seed treated with Onager is to be tagged at processing plants "<b>NOT FOR HUMAN OR ANIMAL CONSUMPTION</b>". It shall be the grower's responsibility to notify the processing plants of any seed crop treated with Onager.</li> <li>• Do not apply more than a total of 24 oz. formulated product per acre per year.</li> <li>• Do not make more than one application per year.</li> </ul>		

- Onager is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager prior to adult mite build up. Onager will not control adult spider mites. Use higher rates on moderate to high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium to high populations, better results may be obtained using Onager in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager.
- Do not make more than one application of Onager or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on the Section 3 label within 120 days of this application.

\* **Types of Irrigation Systems:** Apply Onager only through sprinkler, including center pivot, lateral move, Low Energy Precision Applications (LEPA), end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply Onager through any other type of irrigation system.

**24(c) REGISTRANT**  
Gowan Company  
PO Box 5569  
Yuma, AZ 85366