

## FIFRA §2(ee) RECOMMENDATION

For Distribution and Use only in the State of Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Maine, Maryland, Massachusetts, Michigan, Nebraska, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Vermont, Virginia, Washington, West Virginia and Wisconsin.



EPA Reg. No. 10163-6414

### DIRECTIONS FOR USE

- This recommendation is permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

### FOR MANAGEMENT OF LATE BLIGHT TUBER ROT

When used as part of an integrated late blight management program, according to label instructions, and in conjunction with a sound tuber storage program, Gavel 75DF fungicide will reduce late blight tuber rot incidence in potatoes.

To maximize activity against tuber blight caused by *Phytophthora infestans*, begin applications when conditions favor late blight disease development and in accordance with local late blight advisory recommendations. Continue to make regular applications with Gavel 75DF while these conditions persist and ensure thorough foliar coverage.

Consult the label for Gavel 75DF for recommended treatment schedules and the appropriate pre harvest intervals.