


FENAZAQUIN	GROUP	21A	ACARICIDE
FENAZAQUIN	GROUP	39	FUNGICIDE

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA AND CALIFORNIA**



EPA REG. NO. 10163-322

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE FL OZ/ACRE	RATE LB A.I./ACRE	COMMENTS
LOW GROWING BERRY SUBGROUP 13-07G (1)  Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these	Powdery Mildew	32 -36	0.43 – 0.48	Apply in at least 50 gallons of water per acre. Use higher rates for heavier mite pressure. <ul style="list-style-type: none"> • Do not make more than one application per year. • Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year. • Do not apply this product until crop is through petal fall unless the conditions under the Direction For Use section on the Section 3 label have been met. • PHI = 1 day

RESTRICTIONS ON ALL CROPS:

- Do not apply by any type of irrigation system.
- Only ground sprayer, airblast and aerial applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.
- Do not apply within 100 feet of water bodies (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).
- Do not plant rotational crops, other than those on this label, within 30 days of this application. Do not plant root, tuber and bulb vegetables within 120 days of this application.