


GROUP	21	ACARICIDE
GROUP	39	FUNGICIDE

**FIFRA §2(ee) RECOMMENDATION  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARKANSAS, CALIFORNIA,  
ILLINOIS, INDIANA, MARYLAND, NORTH CAROLINA, SOUTH CAROLINA, PENNSYLVANIA  
AND VIRGINIA.**



**EPA REG. NO. 10163-322**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE FL OZ/ACRE	RATE LB A.I./ACRE	COMMENTS
<b>CANEBERRY SUBGROUP 13-07A</b>  Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and/or hybrids of these	Broad mite	32 -36	0.42 – 0.48	Apply in at least 50 gallons of water per acre. Use higher rates for heavier mite pressure.
				<ul style="list-style-type: none"> <li>• Do not make more than one application per year.</li> <li>• Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year.</li> <li>• Do not apply this product until crop is through petal fall unless the conditions under the Direction For Use section on the Secion 3 label have been met.</li> <li>• Preharvest Interval = 7 days</li> </ul>

**RESTRICTIONS ON ALL CROPS:**

- Aerial applications are prohibited.
- Do not apply by any type of irrigation system.
- Only ground sprayer and airblast applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.
- Do not plant rotational crops, other than those on this label, within 30 days of this application. Do not plant root, tuber and bulb vegetables within 120 days of this application.