

GROUP	10A	ACARICIDE
-------	-----	-----------

**FIFRA §2(ee) RECOMMENDATION FOR ADDITIONAL PEST  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF COLORADO, KANSAS,  
OKLAHOMA AND TEXAS.**



**EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY**

**EPA REG. NO. 10163-277**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE	COMMENTS
<b>CORN, SWEET (including ears and forage) (28)</b> (WESTERN US <u>ONLY</u> *)	Carmine Mite	12 - 24	<b>Ground Application:</b> Apply the recommended rate of Onager with conventional spray boom equipment at a minimum of 20 gallons finished spray per acre. Thorough and uniform coverage of plants is required for optimal results. <b>Air Application:</b> Apply the recommended rate of Onager by air in a minimum of 10 gallons finished spray per acre. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. <b>Chemigation:</b> see details under Chemigation Systems.
			<ul style="list-style-type: none"> <li>• Do not apply more than a total of 24 ozs. formulated product per acre per calendar year.</li> <li>• Do not make more than one application per calendar year.</li> <li>• Do not apply within 28 days of harvest of sweet corn or forage.</li> </ul> *May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281
<b>CORN, FIELD (30)</b>	Carmine Mite	12 - 24	<b>Ground Application:</b> Apply the recommended rate of Onager with conventional spray boom equipment at 15 – 20 GPA. Thorough and uniform coverage of plants is required for optimal results. <b>Air Application:</b> Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. <b>Chemigation:</b> see details under Chemigation Systems.
			<ul style="list-style-type: none"> <li>• Do not apply more than a total of 24 ozs. formulated product per acre per calendar year.</li> <li>• Do not make more than one application per calendar year.</li> <li>• Do not harvest field corn forage, stover, and grain for 30 days after application.</li> </ul>

#### **RESTRICTIONS ON ALL CROPS**

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager prior to adult mite build up. Onager will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using Onager in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager.
- Do not make more than one application of Onager or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.

Onager® and the donkey logo are registered trademarks of Gowan Company, L.L.C.  
19-07-01