

HEXYTHIAZOX	GROUP	10A	ACARICIDE
-------------	-------	-----	-----------

This supplemental label expires on June 20, 2023,  
and must not be used or distributed after this date.

# ONAGER OPTEK MITICIDE

EMULSION FOR AGRICULTURAL USE ONLY

<b>ACTIVE INGREDIENT:</b> Hexythiazox	<b>% By Wt.</b>
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide*	11.93%
<b>OTHER INGREDIENTS**</b>	88.07%
	<b>TOTAL</b> 100.00%

\*Onager Optek contains 1.0 lb. active ingredient per gallon

\*\*Contains Petroleum Distillate

## KEEP OUT OF REACH OF CHILDREN CAUTION

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

### APPLICATION DIRECTIONS

#### PREHARVEST INTERVAL

The required days between the last application and harvest are given in ( ) after each crop name.

CROP	PEST	RATE Oz./acre	INSTRUCTIONS
<b>DATES (90)</b>	Banks Grass Mite	12 - 32	Apply at first signs of egg deposition, but no later than first observation of mite webbing on dates. <b>Ground Application:</b> Apply by directing spray toward bunches and on crown and fronds 3 feet above and below date bunches. Use volume and pressure to assure thorough uniform coverage of all dates. <b>Air Application:</b> apply a minimum of 5 gallons finished spray per acre. <b>Chemigation:</b> see details under Chemigation Systems.
	Restrictions for Dates <ul style="list-style-type: none"> <li>Do not make more than one application per year.</li> <li>Do not apply more than 32 ozs (0.25 lbs ai) of formulated product per acre per year.</li> </ul>		
<b>CANEBERRY SUBGROUP 13-07A:</b> Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and/or hybrids of these (3)	Two-spotted spider mite, McDaniel	12 - 32	<b>Ground Application:</b> Apply a minimum of 20 gallons of finished spray per acre. <b>Air Application:</b> apply a minimum of 5 gallons finished spray per acre. <b>Chemigation:</b> see details under Chemigation Systems.
	Spider Mite; Yellow Spider Mite; Pacific Mite		
Restrictions for Caneberry Subgroup 13-07A <ul style="list-style-type: none"> <li>Do not make more than one application per year.</li> <li>Do not apply more than 32 ozs (0.25 lbs ai) of formulated product per acre per year.</li> <li>Application may be made post harvest</li> </ul>			

### USE RESTRICTIONS ON ALL CROPS

- Apply this product as specified on the EPA approved label and any supplemental labeling.
- Onager Optek is an emulsion in water for application in commercial plantings only.
- Apply Onager Optek prior to adult mite build up. Onager Optek will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using Onager Optek in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager Optek.
- Do not make more than one application of Onager Optek or any other Hexythiazox product per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.

**Gowan**<sup>®</sup>  
The Go To Company