

HEXYTHIAZOX	GROUP	10A	ACARICIDE
-------------	-------	-----	-----------

SUPPLEMENTAL LABELING

This supplemental label expires on August 30, 2022, and must not be used or distributed after this date.

ONAGER OPTEK MITICIDE

EMULSION FOR AGRICULTURAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide*	11.93%
OTHER INGREDIENTS**	<u>88.07%</u>
	TOTAL 100.00%

*Onager Optek contains 1.0 lb. active ingredient per gallon

**Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Onager Optek Miticide before applying. Carefully following all precautionary statements and applicable use directions.
- Use of Onager Optek Miticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Onager Optek Miticide

APPLICATION DIRECTIONS

CROP	PEST	RATE Oz./acre	COMMENTS
LOW GROWING BERRY SUBGROUP 13-07G: Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these (3)	Two-spotted spider mite	12 - 32	Ground Application: Apply a minimum of 20 gallons of finished spray per acre. Air Application: apply a minimum of 20 gallons finished spray per acre. Chemigation: see details under Chemigation Systems.
			<ul style="list-style-type: none"> • Do not make more than one application per year. • Do not apply more than a total of 32 ozs. (0.25 lbs ai) of formulated product per acre per year. • PHI = 3 days

USE RESTRICTIONS ON ALL CROPS

- Apply this product as specified on the EPA approved label and any supplemental labeling.
- Onager Optek is an emulsion to be diluted with water for application in commercial plantings only.
- Apply Onager Optek prior to adult mite build up. Onager Optek will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using Onager Optek in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager Optek.
- Do not make more than one application of Onager Optek or any other hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.


Gowan[®]
The Go To Company

EPA Reg. No. 10163-337
EPA Est. No.

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366