

Group **U6** Fungicide

Supplemental Label



This supplemental label expires on April 30, 2020 and must not be used or distributed after this date.

EPA Reg. No. 8033-103-10163

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Torino Fungicide before applying. Use of Torino Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container of Torino Fungicide.

FRUITING VEGETABLES (within Crop Group 8-10)

SITE	DISEASE	TORINO FUNGICIDE OZ/ACRE [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
FRUITING VEGETABLES: Cocona, Eggplant (including African, Pea and Scarlet), Garden huckleberry, Goji berry, Ground cherry, Martynia, Naranjilla, Okra, Pepino, Peppers (including bell and non-bell), Roselle, Sunberry, Tomatillo, and Tomatoes (including Bush, Currant, and Tree), and cultivars, varieties and/or hybrids of these.	Powdery mildew	3.4 [0.022]	Begin application at first sign of disease development	Do not make more than three (3) applications per year. Do not apply more than once every fourteen (14) days. Do not exceed a total of 0.066 lbs. active ingredient (10.2 oz. of product) per acre per calendar year. Applications may be made up to and including the day of harvest; (PHI = 0 days)