

**REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY IN THE STATE OF WYOMING**
THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION
This label valid until December 31, 2026 or until otherwise amended, withdrawn, canceled or suspended



EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY
EPA Reg. No. 10163-337

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.93%
OTHER INGREDIENTS	88.07%
	TOTAL 100.00%

Onager Optek contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the USEPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

CROP	PEST	RATE OzS./Acre	COMMENTS
ALFALFA GROWN FOR SEED AND CLOVER GROWN FOR SEED	Twospotted Spider Mite, Pacific Spider Mite, Strawberry Spider Mite	12 – 16	Onager Optek is active against all immature stages of Tetranychid spider mites, including freshly laid eggs. Therefore, it is important to apply Onager Optek prior to adult mite buildup. Scout and apply early when mites first begin colonizing the upper 1/3 to 1/2 of the plant canopy. An early application timing and thorough coverage are essential for acceptable control. Apply Onager Optek by ground in a minimum of 20 gallons of water per acre at a minimum of 35 psi to insure optimum coverage.
	<ul style="list-style-type: none"> • Do not cut current year's treated seed crop for hay or forage. • Do not graze current year's treated seed crop. • Screenings from seed processing are prohibited from food or feed channels. All Onager Optek treated seed screenings must be removed from the food or feed market. • Treated seed is not to be used for sprouting. All seed treated with Onager Optek is to be tagged at processing plants "NOT FOR HUMAN OR ANIMAL CONSUMPTION". It shall be the grower's responsibility to notify the processing plants of any seed crop treated with Onager Optek. 		

Restrictions

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager Optek is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager Optek prior to adult mite build up. Onager Optek will not control adult spider mites. Use higher rates on moderate to high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium to high populations, better results may be obtained using Onager Optek in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager Optek.
- Do not make more than one application of Onager Optek or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application
- Do not apply within 4 weeks of harvest

24(c) REGISTRANT

Gowan Company
PO Box 5569
Yuma, AZ 85366