

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
OKLAHOMA AND NEW MEXICO.**



EPA Reg. No. 81880-6-10163

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

FIELD CORN – Reduced Rate Recommendation

For coarse textured soils with an organic matter \leq 1.5% and/or a pH \geq 7.5, apply 3 oz per acre.

YUKON Herbicide may be applied broadcast over-the-top or with drop nozzles to field corn from spike up to 36 inch field corn. To maximize efficacy apply from spike stage through 20 inch field corn. The use of drop nozzles is recommended for applications made corn over 20 inches in height to ensure proper coverage of weeds.

Following application to foliage, corn may be grazed or harvested for feed after the crop reaches the ensilage (milk) stage, at least 30 days after foliar application.

GRAIN SORGHUM- Reduced Rate Recommendation

For coarse textured soils with an organic matter \leq 1.5% and/or a pH \geq 7.5, apply 3 oz per acre.

YUKON Herbicide alone can be applied from the 2-leaf through 15-inch tall sorghum. Use drop nozzles if sorghum is greater than 8 inches. Application made when the sorghum is in the 3 to 5 leaf stage and weeds are small will result in best performance.

Do not graze or feed treated sorghum forage or silage for 30 days after a treatment of YUKON Herbicide. Do not apply to sorghum grown for seed production.