



EPA Reg. No. 10163-331

For use on Potatoes for control of Black dot (*Colletotrichum coccodes*), Botrytis vine rot (*B. cinerea*), Brown spot (*A. Alternaria*), Early blight (*Alternaria solani*), and Late blight (*Phytophthora infestans*) in Washington, Idaho, Oregon, Nebraska, Colorado, Montana, Wisconsin, Michigan, Minnesota, Ohio, North Dakota, and Pennsylvania.

FIFRA §2(ee) RECOMMENDATION: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.
- For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation.
- Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Crop	Diseases Controlled	Rate per Acre (fl oz/Acre)	Remarks (Also Refer to Directions for Use)
Potatoes (7)	Black dot, (<i>Colletotrichum coccodes</i>) Botrytis vine rot, (<i>B. cinerea</i>) Brown spot (<i>A. Alternaria</i>) Early blight, (<i>Alternaria solani</i>) Late blight, (<i>Phytophthora infestans</i>)	24-34 fl oz/A (0.19 lb zoxamide and 1.11 lb chlorothalonil)	Apply on a preventative schedule. Begin applications when conditions are favorable for disease development. Use a 5 to 7 day schedule (the minimum re-treatment interval is 5 days) when disease is present and environmental conditions favor continued disease development. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. Use the maximum labeled rate at row fill. Under low disease conditions and environmental conditions unfavorable for disease development, a 7 to 14 day application schedule may be used. Increase water spray volume as canopy density increases. Apply by ground, aerial or chemigation. Do not exceed a 10 day interval between applications when using chemigation.
	Restrictions		
<ul style="list-style-type: none"> • Do not make more than 2 sequential applications before alternating with a fungicide that has a different mode of action. • Do not make more than 8 applications or apply more than 1.52 lb zoxamide and 8.88 lb chlorothalonil per acre per season. • Do not apply within 7 days of harvest 			

Zing® is a registered trademark used under license of Gowan Company L.L.C.



Produced For :
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569